

linear, logarithm and exponential. Data did not apply to the logistic and the hyperbolic models (Table 3). The best growth model was the power law with the lowest BIC in nearly all species, sex and organs, except for the length of abdomen of male *P. vignaudii* where quadratic function was the best (Table 3). However, the Gompertz model appears as an additional one to the power law model for the elytra in *A. acutipennis*, male *E. p. ibandana*, female with six nymphal instars *E. p. ibandan* and male *P. vignaudii*, as well as for the hind wing in female with six nymphal instars *E. p. ibandan* and male *P. vignaudii* (Table 3).

Growth Ratio

The growth ratio of each organ from one instar to the next varied between stages, sexes and species (Table 4). In all species, the growth ratios of elytra length and hind wing length were high and varied between 2 and 4, except for *E. p. ibandana* F7 where the growth ratios between nymphal instars 6 and 7 were low (between 1 and 1.07). Meanwhile, the growth ratio for the other 13 organs varied between 0.8 and 2.

Table 3. Summary of BIC values for all growth models tested in the organs of *A. acutipennis* (Atrac), *E. p. ibandana* (Eypre), *P. vignaudii* (Pyrgo) and *T. ferruginea* (Taphr).

Species	Sex	Model	Lt	La	Lcc	lcc	Lth	Lpr	Labd	Lel	Laï	Lcu1	Lcu2	Lcu3	Lti1	Lti2	Lti3
Atrac	MF	Exponential	1730	910	963	290	931	833	1254	206	178	655	657	1272	540	547	1190
		Linear	1244	308	392	-77	486	350	920	206	174	78	96	643	0.25	43	563
		Quadratic	1238	292	380	-152	420	241	923	206	140	66	44	587	-36	-5	492
		Power law	-352	-338	-384	-264	-204	-244	-177	19	-16	-292	-371	-416	-318	-295	-416
		Logarithm	1244	308	392	-77	486	350	920	206	174	78	96	643	0.25	43	563
		Gompertz	307	305	228	99	379	434	319	19	56	348	321	321	322	287	304
		Logistic	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
		Hyperbole	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Eypre	M	Exponential	1004	590	490	383	490	437	834	112	75	330	719	845	320	420	716
		Linear	624	93	7	-22	178	-105	544	112	75	-63	373	341	-88	-0.1	219
		Quadratic	618	72	-3.4	-42	183	-116	532	112	75	-63	377	338	-91	-22	216
		Power law	-298	-278	-377	-417	-252	-318	-174	-112	-116	-256	-76	-334	-264	-212	-354
		Logarithm	624	93	7	-22	178	-105	544	112	75	-63	373	341	-88	-0.1	219
		Gompertz	165	167	65	100	92	198	249	-112	-115	133	305	177	125	166	103
		Logistic	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
		Hyperbole	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	F6	Exponential	962	619	533	424	515	483	798	184	52	380	414	855	349	456	749
		Linear	675	393	225	98	243	195	548	184	52	135	156	541	30	114	463
		Quadratic	597	287	157	63	205	56	515	184	52	79	88	410	-129	-9	316
		Power law	-171	-87	-150	-211	-152	-121	-121	-18	-120	-117	-111	-135	-120	-117	-131
		Logarithm	675	393	225	98	243	195	548	184	52	135	156	541	30	114	463
		Gompertz	170	176	100	128	142	214	220	-18	-120	158	158	201	146	189	147
		Logistic	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
		Hyperbole	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	F7	Exponential	1387	930	787	722	822	797	1182	440	414	666	719	1223	617	725	1099
		Linear	980	573	357	393	449	410	782	389	361	289	373	827	209	317	732
		Quadratic	985	561	363	393	455	408	787	151	139	293	378	830	214	322	322
		Power law	-165	-83	-200	-168	-130	-92	-108	136	128	-98	-76	-127	-123	-122	-137
		Logarithm	980	573	357	393	449	410	782	389	361	289	373	827	209	317	732
		Gompertz	308	304	202	258	262	355	362	204	195	302	305	375	280	314	279
		Logistic	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
		Hyperbole	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Pyrgo	M	Exponential	547	603	270	86	459	203	414	46	46	23	34	347	21	17	337
		Linear	365	449	70	-87	290	-5	273	46	46	-0.4	2	113	7	1	144
		Quadratic	366	329	74	-83	227	-80	-147	46	46	-1.5	0.5	115	7	-5	146
		Power law	-115	-79	-89	-131	-75	-98	-69	-30	-22	-38	-37	-144	-23	-34	-112
		Logarithm	365	449	70	-87	290	-5	273	46	46	-0.4	2	113	7	1	144
		Gompertz	102	64	51	36	89	108	115	-30	-22	3	14	78	9.8	-10	78.5
		Logistic	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
		Hyperbole	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	F	Exponential	635	334	337	117	424	254	449	40	35	64	68.5	430	53	50	414
		Linear	436	163	86	-45	305	99	342	45	46	31	40.3	195	9	32	209
		Quadratic	418	117	73	-56	286	60.5	330	29	27	21	13	164	-1.6	5	180
		Power law	-135	-74	-189	-119	-83	-48	-45	-26	-19	-35	-31.5	-151	-39	-23	-128
		Logarithm	436	163	86	-45	305	99	342	45	46	31	40.3	195	9	32	209
		Gompertz	139	117	67	56	461	134	137	-0.4	3	25	33.5	113	38	26	113
Logistic		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
Hyperbole		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
Taphr	M	Exponential	867	558	425	173	424	398	680	198	178	250	281	599	236	547	596
		Linear	640	386	149	-27	305	169	533	160	174	125	158	346	110	43	380
		Quadratic	628	308	125	-42	286	101	526	127	74	76	113	294	68	-5	339
		Power law	-202	-128	-238	-197	-83	-102	-109	-83	-16	-147	-149	-208	-131	-295	-179
		Logarithm	640	386	149	-27	305	169	533	160	174	125	158	346	110	43	380
		Gompertz	23	49	19	-29	46	117	56	67	56	-24	-29	16	-14.3	-33	33.2
		Logistic	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
		Hyperbole	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	F	Exponential	917	603	466	194	459	432	733	228	179	287	315	643	269	286	626
		Linear	701	449	225	23	290	276	573	200	192	184	169	417	190	181	409
		Quadratic	630	329	192	-62	227	157	540	162	122	101	96	332	104	98	328
		Power law	-140	-79	-181	-156	-75	-31	-74	-42	-21	-110	-158	-155	-80	-113	-147
		Logarithm	701	449	225	23	290	276	573	200	192	184	169	417	190	181	409
		Gompertz	44	64	20	-34	57	112	82	60	44	-12	-14	31	-10.3	-17	39
Logistic		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	
Hyperbole		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	

Legend: M: male, F: female, F6: female with six nymphal instars, F7: female with seven nymphal instars, body length (Lt), length (Lcc) and width (lcc) of cephalic capsule, length of thorax (Lth), abdomen (Labd), pronotum (Lpr), antenna (La), elytra (LeI), hind wing (Lai), anterior (Lcu1), median (Lcu2) and posterior (Lcu3) femur, anterior (Lti1), median (Lti2) and posterior (Lti3) tibia. NA: not applicable.

Table 4. Growth ratio (Y_{n+1} / Y_n) of each organ of *A. acutipennis* (Atrmph), *E. p. ibandana* (E.p.i), *P. vignaudii* (Pygmph) and *T. ferruginea* (Tphrn) in the laboratory.

Organs	Stages	Species/Sexes							
		Atrmph	E.p.i-M	E.p.i-F6	E.p.i-F7	Pygmph-M	Pygmph-F	Tphrn-M	Tphrn-F
Lt	L2/L1	1.62±0.04a	1.53±0.06a	1.54±0.06a	1.54±0.06a	1.57±0.1a	1.87±0.07a	1.3±0.05a	1.3±0.06ab
	L3/L2	1.25±0.03b	1.44±0.03a	1.36±0.02b	1.36±0.02b	1.4±0.1a	1.37±0.06b	1.2±0.03ab	1.2±0.04b
	L4/L3	1.3±0.03b	1.11±0.02b	1.16±0.03b	1.16±0.03b	1.2±0.07b	1.14±0.06b	1.13±0.02b	1.16±0.03b
	L5/L4	1.08±0.04b	1.23±0.04b	1.47±0.03a	1.37±0.05b	1.25±0.06b	1.3±0.05b	1.22±0.04ab	1.27±0.05ab
	L6/L5	1.6±0.04a	1.2±0.03b	1.45±0.02a	1.62±0.04a	1.3±0.04b	1.4±0.05b	1.26±0.04ab	1.4±0.05a
	L7/L6				1±0.02c				
	A/L6	1±0.02b	1.42±0.02a	1.42±0.03a		1.06±0.04b	1.4±0.07b	1.2±0.04ab	1.45±0.03a
	A/L7				1.4±0.05a				
Lcc	L2/L1	1.45±0.04a	1.16±0.03a	1.2±0.02a	1.2±0.02a	1.2±0.1a	1.32±0.05a	1.2±0.04ab	1.4±0.04a
	L3/L2	1.35±0.03a	1.4±0.02b	1.3±0.02ab	1.3±0.02b	1.54±0.24b	1.32±0.25a	1.3±0.04b	1.13±0.04b
	L4/L3	1.24±0.02b	1.2±0.02a	1.2±0.02a	1.2±0.02a	1.12±0.03a	1.2±0.25ab	1.1±0.03a	1.16±0.03b
	L5/L4	1.11±0.03b	1.16±0.02a	1.4±0.03b	1.27±0.04ab	1.22±0.04a	1.2±0.03ab	1.2±0.02ab	1.17±0.03b
	L6/L5	1.45±0.04a	1.14±0.02a	1.33±0.03ab	1.47±0.03b	1.15±0.02a	1.3±0.05a	1.3±0.03b	1.33±0.02a
	L7/L6				1.03±0.01c				
	A/L6	1±0.02b	1.25±0.03a	1.2±0.02a		1.07±0.02a	1.17±0.03b	1.2±0.03ab	1.32±0.03a
	A/L7				1.16±0.01a				
lcc	L2/L1	1.14±0.01a	1.47±0.03a	1.53±0.05a	1.53±0.05a	1.25±0.07ab	1.36±0.04a	1.25±0.04a	1.3±0.04a
	L3/L2	1.22±0.02a	1.2±0.02b	1.14±0.01b	1.14±0.01b	1.3±0.05b	1.27±0.3ab	1.2±0.03ab	1.1±0.04b
	L4/L3	1.3±0.07ab	1.23±0.02b	1.2±0.03b	1.2±0.03b	1.2±0.04a	1.17±0.03b	1.12±0.01b	1.16±0.03b
	L5/L4	1.17±0.05a	1.15±0.01b	1.47±0.03a	1.24±0.04b	1.17±0.06a	1.25±0.03ab	1.15±0.02b	1.2±0.03ab
	L6/L5	1.43±0.05b	1.16±0.02b	1.27±0.03ab	1.7±0.05a	1.17±0.03a	1.28±0.07ab	1.25±0.04a	1.33±0.03a
	L7/L6				1±0.02b				
	A/L6	1.1±0.02a	1.12±0.01b	1.15±0.02b		1.2±0.03a	1.36±0.07a	1.2±0.03ab	1.3±0.03a
	A/L7				1.1±0.01b				
Lth	L2/L1	1.46±0.04a	1.3±0.04a	1.5±0.1a	1.5±0.1a	2.2±0.12a	1.86±0.2a	1.35±0.08a	1.35±0.04a
	L3/L2	1.34±0.04a	1.45±0.04b	1.35±0.02ab	1.35±0.02ab	1.18±0.05b	1.82±0.2a	1.23±0.04ab	1.15±0.03b
	L4/L3	1.4±0.03a	1.1±0.02c	1.2±0.02b	1.2±0.02b	1.17±0.06b	1.17±0.03b	1.14±0.02b	1.11±0.04b
	L5/L4	1.27±0.04a	1.16±0.03ac	1.28±0.02b	1.34±0.02ab	1.23±0.06b	1.47±0.1ab	1.22±0.04b	1.4±0.07a
	L6/L5	1.75±0.06a	1.27±0.03ac	1.44±0.02a	1.4±0.01a	1.32±0.06a	1.36±0.1ab	1.36±0.04a	1.32±0.05a
	L7/L6				1.03±0.02c				
	A/L6	1.1±0.01b	1.44±0.04b	1.47±0.03a		1.12±0.04b	1.28±0.05b	1.4±0.05a	1.61±0.05a
	A/L7				1.44±0.03a				
Labd	L2/L1	1.63±0.06a	1.91±0.1a	1.75±0.07a	1.75±0.07a	1.5±0.09a	1.6±0.05a	1.42±0.06a	1.42±0.06a
	L3/L2	1.21±0.04b	1.43±0.04b	1.44±0.04a	1.44±0.04a	1.46±0.08a	1.51±0.07a	1.15±0.04b	1.2±0.05b
	L4/L3	1.24±0.04b	1.12±0.03c	1.14±0.04b	1.14±0.04b	1.36±0.08ab	1.21±0.04b	1.2±0.05b	1.21±0.04b
	L5/L4	1.17±0.03b	1.26±0.05c	1.53±0.04a	1.5±0.07a	1.24±0.07b	1.21±0.04b	1.23±0.06b	1.25±0.07b
	L6/L5	1.27±0.05b	1.24±0.05c	1.43±0.04a	1.6±0.04a	1.3±0.05ab	1.6±0.1a	1.24±0.06b	1.46±0.08a
	L7/L6				1.02±0.02b				
	A/L6	1.23±0.04b	1.35±0.04b	1.55±0.05a		1.16±0.04b	1.63±0.15a	1.25±0.06b	1.57±0.05a
	A/L7				1.45±0.07a				

Lpr	L2/L1	1.6±0.07a	1.35±0.03ab	1.42±0.03ab	1.42±0.03a	1.43±0.08a	1.43±0.1a	1.42±0.1a	1.38±0.06a
	L3/L2	1.42±0.04b	1.5±0.02a	1.42±0.03ab	1.42±0.03a	1.47±0.08a	1.56±0.04a	1.36±0.04a	1.17±0.05b
	L4/L3	1.4±0.03b	1.22±0.03b	1.22±0.03b	1.22±0.03b	1.35±0.07a	1.18±0.06b	1.04±0.02b	1.2±0.06b
	L5/L4	1.27±0.06b	1.35±0.03ab	1.7±0.05a	1.33±0.05ab	1.24±0.07ab	1.44±0.06a	1.48±0.05a	1.5±0.05a
	L6/L5	1.64±0.05a	1.25±0.02b	1.45±0.03a	1.92±0.02a	1.3±0.03ab	1.43±0.05a	1.4±0.04a	1.45±0.03a
	L7/L6				1±0.01b				
	A/L6	0.8±0.01c	1.21±0.02b	1.1±0.02b		1.1±0.01b	1.33±0.06ab	1.21±0.06ab	1.36±0.04ab
	A/L7				1.05±0.02b				
La	L2/L1	1.35±0.04a	1.32±0.04a	1.32±0.03ab	1.32±0.03a	1.35±0.05a	1.46±0.1a	1.27±0.05ab	1.27±0.06ab
	L3/L2	1.36±0.03a	1.41±0.02a	1.34±0.02ab	1.34±0.02a	1.33±0.05a	1.36±0.06ab	1.3±0.04b	1.3±0.06ab
	L4/L3	1.4±0.03a	1.24±0.02a	1.25±0.03b	1.25±0.03a	1.32±0.05a	1.21±0.06b	1.06±0.03a	1.12±0.06b
	L5/L4	1.16±0.04b	1.3±0.02a	1.6±0.06a	1.23±0.05a	1.29±0.05a	1.38±0.1ab	1.36±0.04b	1.4±0.06a
	L6/L5	1.35±0.04a	1.26±0.02a	1.47±0.04a	1.82±0.05b	1.32±0.05a	1.44±0.1a	1.4±0.03b	1.44±0.02a
	L7/L6				1.08±0.03a				
	A/L6	1.22±0.03ab	1.62±0.02b	1.22±0.02b		1.33±0.07a	1.3±0.05ab	1.3±0.03b	1.3±0.04ab
	A/L7				1.16±0.03a				
Lel	L2/L1								
	L3/L2								
	L4/L3								
	L5/L4								
	L6/L5	2.32±0.15a	2.58±0.07a	2.9±0.13a		2.1±0.07a	2.27±0.08a	2.24±0.06a	2.2±0.08a
	L7/L6				1.01±0.02a				
	A/L6	2.86±0.05b	3.76±0.13b	3.7±0.2b		3.1±0.1b	2.7±0.1b	3.74±0.14b	4.4±0.12b
	A/L7				3.43±0.1b				
Lai	L2/L1								
	L3/L2								
	L4/L3								
	L5/L4								
	L6/L5	2.41±0.18a	2.12±0.06a	2.29±0.04a		2.28±0.06a	2.61±0.11a	2.45±0.14a	2.47±0.14a
	L7/L6				1.02±0.02a				
	A/L6	2.46±0.17a	3.74±0.08b	3.77±0.07b		2.9±0.1b	2.21±0.08b	4±0.15b	4.5±0.12b
	A/L7				3.6±0.1b				
Lcu1	L2/L1	1.34±0.03a	1.15±0.01a	1.2±0.03a	1.2±0.03a		1.7±0.12a	1.3±0.09a	1.32±0.1a
	L3/L2	1.5±0.04b	1.51±0.02b	1.47±0.03b	1.47±0.03b		1.4±0.11a	1.16±0.02ab	1.15±0.05b
	L4/L3	1.27±0.03a	1.2±0.02a	1.2±0.03a	1.2±0.03a		1.05±0.13b	1.03±0.02b	1.05±0.04b
	L5/L4	1.45±0.11b	1.25±0.04a	1.52±0.03b	1.3±0.04ab	1.13±0.05a	1.3±0.06a	1.27±0.04a	1.27±0.03ab
	L6/L5	2.28±0.15c	1.26±0.04a	1.4±0.04b	1.77±0.04b	1.36±0.05b	1.5±0.1a	1.34±0.04a	1.44±0.04a
	L7/L6				1.07±0.02a				
	A/L6	2.85±0.05d	1.6±0.03b	1.3±0.04ab		1.24±0.04ab	1.3±0.06a	1.24±0.04a	1.4±0.04a
	A/L7				1.3±0.03a				

Lcu2	L2/L1	1.34±0.03a	1.2±0.03a	1.3±0.04a	1.3±0.04a		1.7±0.06a	1.3±0.07a	1.27±0.06ab
	L3/L2	1.41±0.04a	1.47±0.03b	1.4±0.04a	1.4±0.04a		1.4±0.12a	1.11±0.02b	1.18±0.05a
	L4/L3	1.3±0.04a	1.2±0.04a	1.1±0.03b	1.1±0.03b		1.02±0.12b	1.04±0.02b	1.1±0.05a
	L5/L4	1.23±0.05b	1.3±0.03c	1.7±0.05c	1.35±0.05a		1.3±0.08a	1.3±0.04a	1.23±0.04a
	L6/L5	1.5±0.05a	1.3±0.02c	1.35±0.03a	1.73±0.06c	1.4±0.02a	1.53±0.09a	1.34±0.04a	1.43±0.03b
	L7/L6				1.02±0.04b				
	A/L6	1±0.23b	1.54±0.03b	1.35±0.03a		1.24±0.04b	1.33±0.06a	1.2±0.04ab	1.35±0.05b
	A/L7				1.25±0.03a				
Lcu3	L2/L1	1.45±0.02a	1.33±0.01ab	1.41±0.02a	1.41±0.02a	1.32±0.06ab	1.41±0.03a	1.24±0.04a	1.27±0.05ab
	L3/L2	1.4±0.02a	1.4±0.02b	1.35±0.02b	1.35±0.02b	1.4±0.1a	1.46±0.05a	1.29±0.03a	1.2±0.05ab
	L4/L3	1.28±0.03b	1.3±0.03a	1.25±0.04b	1.25±0.04b	1.26±0.06ab	1.16±0.04b	1.04±0.02b	1.15±0.06a
	L5/L4	1.18±0.03b	1.29±0.03a	1.7±0.05a	1.32±0.05b	1.18±0.05b	1.32±0.04a	1.34±0.02a	1.31±0.03b
	L6/L5	1.5±0.04a	1.25±0.02a	1.37±0.02ab	1.77±0.04c	1.23±0.03ab	1.37±0.05a	1.29±0.02a	1.3±0.02b
	L7/L6				1.01±0.01d				
	A/L6	1±0.01c	1.25±0.02a	1.26±0.03b		1.24±0.03ab	1.3±0.05a	1.24±0.03a	1.4±0.03b
	A/L7				1.25±0.03b				
Lti1	L2/L1	1.34±0.02a	1.15±0.02a	1.2±0.02a	1.2±0.02a		1.9±0.14a	1.3±0.09a	1.25±0.1a
	L3/L2	1.4±0.04a	1.45±0.03b	1.4±0.02b	1.4±0.02b		1.3±0.14b	1.2±0.03a	1.2±0.1a
	L4/L3	1.37±0.03a	1.23±0.03b	1.2±0.03a	1.2±0.03a		1.1±0.08c	1±0.02b	1.08±0.04b
	L5/L4	1.13±0.04b	1.26±0.03b	1.56±0.03b	1.3±0.04ab		1.36±0.05b	1.32±0.05a	1.27±0.03a
	L6/L5	1.53±0.05a	1.24±0.03b	1.37±0.02ab	1.7±0.04c	1.37±0.01a	1.43±0.06b	1.33±0.04a	1.48±0.05a
	L7/L6				1±0.01d				
	A/L6	1±0.2b	1.28±0.03b	1.21±0.01a		1.3±0.06a	1.3±0.04b	1.24±0.03a	1.32±0.04a
	A/L7				1.26±0.02a				
Lti2	L2/L1	1.31±0.02a	1.21±0.03a	1.3±0.03a	1.3±0.03a		1.83±0.08a	1.26±0.06a	1.31±0.06ab
	L3/L2	1.34±0.04a	1.45±0.03b	1.4±0.03a	1.4±0.03a		1.34±0.11a	1.16±0.03a	1.11±0.05a
	L4/L3	1.35±0.04a	1.21±0.03a	1.22±0.03a	1.22±0.03a		1.07±0.12b	1±0.03b	1.2±0.05a
	L5/L4	1.3±0.08a	1.4±0.04b	1.71±0.05b	1.34±0.05a		1.41±0.1a	1.31±0.05a	1.24±0.04a
	L6/L5	1.42±0.06a	1.25±0.02a	1.32±0.02a	1.72±0.05b	1.4±0.03a	1.5±0.1a	1.33±0.04a	1.45±0.04b
	L7/L6				1±0.01c				
	A/L6	1±0.02b	1.29±0.02ab	1.3±0.02a		1.21±0.05b	1.4±0.1a	1.3±0.03a	1.36±0.04b
	A/L7				1.26±0.03a				
Lti3	L2/L1	1.45±0.02a	1.28±0.02a	1.33±0.02a	1.33±0.02a	1.31±0.1ab	1.34±0.02a	1.26±0.06a	1.25±0.05ab
	L3/L2	1.36±0.02ab	1.25±0.02a	1.22±0.01a	1.22±0.01a	1.46±0.1a	1.52±0.05a	1.3±0.03a	1.27±0.05a
	L4/L3	1.28±0.03b	1.26±0.02a	1.26±0.03a	1.26±0.03a	1.22±0.04b	1.11±0.05b	1.05±0.02b	1.15±0.06b
	L5/L4	1.23±0.04b	1.26±0.02a	1.6±0.04b	1.26±0.02a	1.2±0.06b	1.34±0.04a	1.31±0.03a	1.3±0.04a
	L6/L5	1.45±0.04a	1.19±0.02a	1.4±0.02ab	1.73±0.04b	1.24±0.04b	1.4±0.06a	1.32±0.03a	1.36±0.02a
	L7/L6				1±0.01c				
	A/L6	1.28±0.02b	1.4±0.02b	1.23±0.02a		1.15±0.08b	1.28±0.05a	1.25±0.03a	1.38±0.04a
	A/L7				1.35±0.02a				

Notes: Each value of the table represents: mean and standard error. Means were compared by performing One-Way ANOVA test and Tukey post hoc test at the $\alpha = 0.05$ level of significance. Within columns, means with same letters are not significantly different. Legend: La: length of antenna. Labd: length of abdomen. Lai: length of the hind wing. Lcc: length of cephalic capsule. lcc: width of cephalic capsule. Lcu1, Lcu2 and Lcu3: length of femur 1, length of femur 2, length of femur 3. Lel: length of elytra. Lpr: length of pronotum. Lt: length of body. Lti1, Lti2, Lti3: length of tibia 1, length of tibia 2, length of tibia 3. Lth: length of thorax. M: male. F: female. F6: female with six nymphal instars. F7: female with seven nymphal instars. L: nymphal instars. A: adult instars.

Allometric Growth

The allometric growth was not harmonious in the studied grasshopper species (Table 5). For each organ, there is a disproportionate variation of the allometric coefficient from one instar to another. Hypoallometries were the most dominant in all four grasshopper species. However, 7 cases of hyperallometries were noticed namely in the width of cephalic capsule

in the adult instar of *E. p. ibandana* M ($k=1.5\pm 1.6$, $p<0,05$); the Pronotum's length in *E. p. ibandana* F7 (in the fifth ($k=1.1\pm 0.08$, $p<0,05$) and the seventh ($k=1.14\pm 1.16$, $p<0,05$) nymphal instars; the lengths of tibia 1 and 2 in *E. p. ibandana* F6 and *E. p. ibandana* F7, in the sixth nymphal instars ($k=1.43\pm 0.4$ for tibia 1 and $k=4.1\pm 0.36$ for tibia 2 (the strongest one), $p<0,05$) and the seventh nymphal instars

Table 5. Allometric coefficient (k) of each organ of *A. acutipennis* (Atrmph), *E. p. ibandana* (E.p.i), *P. vignaudii* (Pygmph) and *T. ferruginea* (Tphrn) in the laboratory.

Organs	Stages	Species/Sexes							
		Atrmph	E.p.i-M	E.p.i-F6	E.p.i-F7	Pygmph-M	Pygmph-F	Tphrn-M	Tphrn-F
Lcc	L1	-0.6±0.1*	-0.9±0.3*	-1.3±0.4*	-1.3±0.4*	-0.64±0.3*	-0.7±0.2*	-0.9±0.3*	-0.5±0.3*
	L2	-0.8±0.1*	-0.5±0.3*	-0.6±0.1*	-0.6±0.1*	-0.55±0.2*	-0.4±0.4*	-0.65±0.2*	-0.3±0.2*
	L3	-0.5±0.1*	-0.9±0.3*	-1.1±0.3*	-1.1±0.3*	0.14±0.4*	-0.2±0.3*	-0.7±0.2*	-0.7±0.1*
	L4	-0.6±0.2*	-0.4±0.2*	-0.15±0.15*	-0.15±0.15*	0.4±0.3*	-0.3±0.3*	-0.18±0.18*	-0.7±0.1*
	L5	0.1±0.1*	-1.2±0.3*	-0.5±0.1*	0.02±0.07*	-0.7±0.7*	-0.2±0.2*	-0.4±0.2*	-0.3±0.2*
	L6	-0.3±0.2*	-0.5±0.4*	-0.75±0.2*	-0.54±0.6*	-0.7±0.3*	0.4±0.3*	-0.04±0.3*	-0.4±0.4*
	L7				-1.3±0.4*				
	Adult	0.3±0.07*	-0.75±0.1*	-0.6±0.2*	-0.2±0.7*	0.03±0.3*	0.2±0.4*	-0.2±0.2*	-0.8±0.1*
lcc	L1	-0.72±0.7*	-0.5±0.2*	-0.56±0.2*	-0.56±0.2*	-1.15±0.2*	-0.9±0.16*	-0.62±0.2*	-0.6±0.23*
	L2	-0.54±0.27*	-0.5±0.3*	-0.56±0.44*	-0.56±0.44*	0.04±0.36*	-0.7±0.4*	-0.99±0.2*	-0.2±0.2*
	L3	-1.2±0.1*	-0.96±0.33*	0.45±0.4*	0.45±0.4*	-0.7±0.3*	-1±0.24*	-0.9±0.22*	-0.56±0.1*
	L4	-1.14±0.1*	-0.52±0.1*	-0.36±0.16*	-0.36±0.16*	-0.5±0.3*	-0.85±0.3*	-0.42±0.3*	-0.45±0.18*
	L5	-0.7±0.2*	-0.06±0.5*	0.08±0.1*	0.25±0.1*	-1.3±0.4*	-0.7±0.2*	-0.36±0.16*	-0.24±0.2*
	L6	-0.72±0.2*	-0.6±0.16*	-0.24±0.07*	-0.8±0.02*	-0.95±0.3*	-0.4±0.2*	-0.06±0.16*	-0.4±0.2*
	L7				0.23±0.1*				
	Adult	-0.3±0.07*	1.5±1.6*	-0.36±0.2*	-0.32±0.33*	-1.2±0.2*	-1±0.3*	-0.24±0.2*	-0.9±0.12*
Lth	L1	-0.7±0.07*	-0.5±0.1*	-0.55±0.1*	-0.55±0.1*	-0.75±0.2*	-0.76±0.07*	-0.55±0.1*	-0.53±0.14*
	L2	-0.65±0.07*	-0.4±0.1*	-0.62±0.16*	-0.62±0.16*	-0.33±0.14*	-0.64±0.12*	-0.62±0.1*	-0.4±0.1*
	L3	-0.84±0.1*	-0.3±0.1*	-0.41±0.1*	-0.41±0.1*	0.02±0.24*	-0.11±0.12*	-0.62±0.12*	-0.7±0.1*
	L4	-0.4±0.1*	-0.27±0.1*	-0.2±0.1*	-0.2±0.1*	-0.04±0.4*	-0.56±0.2*	-0.9±0.1*	-0.72±0.1*
	L5	-0.5±0.1*	-0.2±0.1*	-0.3±0.16*	-0.54±0.05*	-0.61±0.2*	-0.8±0.07*	-0.5±0.1*	-0.6±0.2*
	L6	-0.4±0.1*	-0.5±0.16*	-0.07±0.06*	-0.5±0.06*	-0.61±0.2*	-0.35±0.24*	-0.5±0.13*	-0.7±0.16*
	L7				0.14±0.1*				
	Adult	-0.3±0.06*	-0.86±0.1*	-0.25±0.1*	-0.21±0.2*	-0.23±0.23*	-0.6±0.25*	-0.42±0.12*	-0.84±0.07*
Labd	L1	-0.55±0.08*	*0.42±0.13*	-0.6±0.1*	-0.6±0.1*	-0.7±0.13*	-0.75±0.08*	-0.36±0.12*	-0.52±0.08*
	L2	-0.5±0.05*	-0.36±0.1*	-0.4±0.1*	-0.4±0.1*	-0.1±0.1*	-0.5±0.1*	-0.34±0.08*	-0.4±0.12*
	L3	-0.34±0.05*	-0.3±0.06*	-0.3±0.06*	-0.3±0.06*	-0.3±0.1*	-0.34±0.1*	-0.44±0.07	-0.54±0.12*
	L4	-0.42±0.04*	-0.4±0.07*	-0.07±0.07*	-0.07±0.07*	-0.31±0.12*	-0.4±0.1*	-0.62±0.1*	-0.5±0.05*
	L5	-0.4±0.1*	-0.35±0.03*	-0.46±0.15*	-0.1±0.01*	-0.55±0.16	-0.8±0.2*	-0.6±0.08*	-0.5±0.05*
	L6	-0.5±0.07*	-0.4±0.03*	-0.75±0.1*	-0.3±0.08*	-0.5±0.1*	-0.3±0.1*	-0.45±0.07*	-0.52±0.08*
	L7				-0.2±0.02*				
	Adult	-0.24±0.03*	-0.6±0.2*	-0.44±0.05*	-0.43±0.1*	-0.74±0.11*	-0.4±0.03*	-0.36±0.06*	-0.6±0.05*

Lpr	L1	-0.64±0.08*	-0.25±0.37*	-1.4±0.6*	-1.4±0.6*	-0.9±0.16*	-1.1±0.08*	-0.75±0.12*	-0.66±0.14*
	L2	-1.2±0.1*	-0.3±0.2*	-0.7±0.2*	-0.7±0.2*	-0.11±0.2*	-0.9±0.33*	-0.73±0.17*	-0.6±0.14*
	L3	-0.55±0.13*	-0.3±0.3*	-0.1±0.5*	-0.1±0.5*	-0.25±0.26*	-0.44±0.3*	-1.03±0.17*	-0.7±0.1*
	L4	-0.52±0.12*	-0.42±0.12*	-0.4±0.1*	-0.4±0.1*	-0.3±0.3*	-0.7±0.2*	-0.93±0.3*	-0.82±0.1*
	L5	-0.2±0.06*	0.24±0.7*	-0.12±0.24*	1.1±0.08*	-1.05±0.5*	-0.46±0.16*	-0.5±0.2*	-0.9±0.14*
	L6	-0.5±0.3*	-0.6±0.22*	-0.13±0.13*	-0.4±0.08*	-0.6±0.25*	-0.43±0.25*	-0.25±0.22*	-0.41±0.22*
	L7				1.14±0.16*				
	Adult	-0.2±0.05*	-0.7±0.16*	-0.42±0.15*	0.11±0.25*	-0.51±0.3*	-0.43±0.4*	-0.63±0.1*	-0.81±0.1*
La	L1	-0.7±0.1*	-0.62±0.15*	0.07±0.23*	0.07±0.23*	-1.07±0.14*	-0.64±0.13*	-0.61±0.2*	-0.65±0.23*
	L2	-0.6±0.2*	-0.5±0.33*	-0.92±0.23*	-0.92±0.23*	-0.06±0.36*	-0.24±0.2*	-0.36±0.2*	-0.3±0.2*
	L3	-0.9±0.07*	-1.3±0.44	-0.53±0.35*	-0.53±0.35*	-0.28±0.27*	-0.61±0.22*	-1.02±0.14*	-0.7±0.1*
	L4	-0.6±0.14*	-0.08±0.16*	-0.4±0.13	-0.4±0.13	-0.44±0.34*	-0.77±0.2*	-0.76±0.21*	-0.75±0.1*
	L5	-0.32±0.18*	-0.8±0.34*	-0.9±0.11*	-0.5±0.12*	-0.64±0.4*	-0.63±0.2*	-0.91±0.25*	-0.7±0.23*
	L6	-0.23±0.16*	-0.6±0.2*	-0.5±0.1*	-0.9±0.03*	-0.8±0.23*	-0.65±0.22*	-0.66±0.23*	-0.5±0.22
	L7				-0.3±0.07*				
	Adult	-0.76±0.3*	-1.83±4.9*	-0.72±0.2*	-0.06±0.4*	-0.6±0.2*	-0.35±0.3*	-0.52±0.32*	-0.84±0.1*
Lel	L1								
	L2								
	L3								
	L4								
	L5	-0.73±0.08*	-0.52±0.31*	-0.8±0.11*	-1.34±0.05	-0.81±0.35*	-1.1±0.22*	-0.98±0.1*	-0.8±0.08*
	L6	-0.7±0.22*	-0.7±0.1*	-0.91±0.06*	-0.4±0.13*	-0.83±0.12*	-0.5±0.35*	-0.84±0.17*	-0.53±0.1*
	L7				-0.41±0.1*				
	Adult	-0.05±0.05*	-1.03±0.02*	-0.51±0.17*	-0.01±0.18*	-0.85±0.17*	-0.84±0.3*	-0.43±0.1*	-0.92±0.06*
Lai	L1								
	L2								
	L3								
	L4								
	L5	-0.8±0.08	-0.63±0.2*	-0.5±0.11*	/	-0.91±0.3*	-0.99±0.27*	-1.05±0.07*	-0.79±0.08*
	L6	-0.6±0.2*	-0.65±0.12*	0.21±0.07*	-0.8±0.02*	-1±0.08*	-1.3±0.37*	-0.8±0.11*	-0.72±0.14*
	L7				0.01±0.09*				
	Adult	0.04±0.07*	-1.03±0.02*	-0.6±0.14*	-0.15±0.16*	-0.8±0.17	-1.15±0.25*	-0.42±0.1*	-0.93±0.06*
Lcu1	L1	-0.72±0.08*	-1.73±0.55*	-0.96±0.7*	-0.96±0.7*	±	±	-0.66±0.14*	-0.67±0.15*
	L2	-0.92±0.11*	-0.99±0.4*	-1.05±0.22*	-1.05±0.22*	±	-0.82±0.35*	-0.85±0.25*	-0.01±0.19*
	L3	-0.78±0.12*	-0.51±0.24*	-1.17±0.21*	-1.17±0.21*	±	-0.25±0.04*	-0.64±0.21*	-0.72±0.1*
	L4	-0.73±0.22*	-0.41±0.21*	-0.32±0.18*	-0.32±0.18*	±	-0.55±0.2*	-0.65±0.17*	-0.62±0.14*
	L5	-0.14±0.12*	-0.59±0.11*	-0.3±0.14*	-0.11±0.13*	-0.87±0.37*	-0.55±0.15*	-0.32±0.18*	-0.7±0.24*
	L6	-0.44±0.14*	-0.77±0.22*	-0.93±0.1*±	-1.36±0.11	-0.92±0.2*	-0.85±0.23*	-0.47±0.17*	0.55±0.14*
	L7				-0.5±0.16*				
	Adult	0±0.11*	-0.72±0.16*	-0.55±0.17*	-0.25±0.33*	-0.85±0.3*	-0.88±0.25*	-0.46±0.21*	-0.81±0.08*

Lcu2	L1	-0.88±0.04*	-0.94±0.43*	-1.27±0.5*	-1.27±0.5*	±	-1.8±0.4*	-0.63±0.2*	-0.62±0.27*
	L2	-0.93±0.07*	-1.47±0.27*	-1.02±0.16*	-1.02±0.16*	-0.6±0.2*	-2.3±0.7*	-0.76±0.25*	-0.03±0.25*
	L3	-0.82±0.16*	-0.9±0.13*	-0.98±0.27*	-0.98±0.27*	0.2±1.45*	-0.53±0.18*	-0.96±0.14*	-0.66±0.12*
	L4	-1.05±0.2*	-0.7±0.15*	-0.4±0.12*	-0.4±0.12*	±	-0.56±0.17*	-0.8±0.17*	-0.68±0.11*
	L5	-0.28±0.14*	-0.35±0.28*	-0.68±0.11*	-0.1±0.1*	±	-0.53±0.17*	-0.5±0.17*	-0.26±0.21*
	L6	-0.17±0.22*	-0.7±0.15*	-0.46±0.11*	-0.43±0.02*	-1.03±0.22*	-1.13±0.27*	-0.55±0.18*	-0.46±0.15*
	L7				-0.02±0.05*				
	Adult	0.2±0.1*	-0.67±0.18*	-0.55±0.18*	-0.13±0.26*	-0.65±0.24*	-0.38±0.28*	-0.37±0.2*	-0.83±0.07*
Lcu3	L1	-0.47±0.09*	0±0.7*	0.997±0.7	0.997±0.7	-0.37±0.42*	-0.52±0.29*	-0.4±0.26*	-0.41±0.22*
	L2	-0.94±0.07*	-0.02±0.4*	-0.46±0.3*	-0.46±0.3*	-0.11±0.19*	0.56±0.37*	-0.56±0.22*	-0.44±0.27*
	L3	-0.44±0.14*	-0.4±0.2*	-0.83±0.35*	-0.83±0.35*	-0.17±0.39*	-0.65±0.3*	-0.85±0.21*	-0.72±0.1*
	L4	-0.4±0.2*	-0.31±0.15*	-0.31±0.12*	-0.31±0.12*	-0.14±0.33*	-0.62±0.21*	-0.64±0.31*	-0.7±0.12*
	L5	-0.46±0.15*	-0.4±0.2*	-0.39±0.13*	0.35±0.18	-0.31±0.7*	-0.08±0.22*	-0.51±0.26*	-0.33±0.36*
	L6	-0.2±0.24*	-0.54±0.2*	-0.01±0.17*	-0.71±0.05*	-0.75±0.36*	-0.04±0.33*	-0.08±0.28*	-0.29±0.28*
	L7				-0.22±0.17*				
	Adult	-0.16±0.07*	-0.23±0.15*	-0.72±0.12*	-0.16±0.23*	-0.33±0.36*	-0.3±0.4*	-0.23±0.22*	-0.77±0.11*
Lti1	L1	-0.61±0.1*	-0.43±0.67*	-0.57±0.73*	-0.57±0.73*	±	±	-0.77±0.14*	-0.72±0.16*
	L2	-0.89±0.13*	-1.2±0.4*	-1.1±0.24*	-1.1±0.24*	±	±	-0.61±0.2*	-0.3±0.15*
	L3	-0.88±0.12*	-0.95±0.15*	-1.22±0.2*	-1.22±0.2*	±	0.11±0.43*	-0.54±0.25*	-0.75±0.1*
	L4	-0.72±0.18*	-0.7±0.17*	-0.35±0.2*	-0.35±0.2*	±	-0.51±0.22*	-0.81±0.15*	-0.78±0.11*
	L5	-0.43±0.05*	-1.15±0.2*	-0.67±0.13*	-0.38±0.27*	-0.94±0.34*	-0.47±0.1*	-0.36±0.014*	-0.78±0.23*
	L6	-0.88±0.2*	-0.7±0.17*	1.43±0.4*	-0.51±0.07*	-1.2±0.15*	-0.5±0.37*	-0.7±0.21*	-0.64±0.14*
	L7				1.66±0.53*				
	Adult	0.26±0.1*	-0.7±0.18*	-0.43±0.17*	0.01±0.32	-0.7±0.5*	-1.12±0.29*	-0.6±0.18*	-0.87±0.07*
Lti2	L1	-0.57±0.07*	-1.3±0.21*	-0.27±0.47*	-0.27±0.47*	±	±	-0.71±0.2*	-0.64±0.22*
	L2	-0.97±0.07*	-1.22±0.26*	-1.03±0.2*	-1.03±0.2*	±	-1.25±0.7*	-0.8±0.25*	-0.23±0.27*
	L3	-0.89±0.11*	-0.85±0.18*	-0.7±0.24*	-0.7±0.24*	-0.36±0.65*	-0.42±0.52*	-0.95±0.15*	-0.78±0.1*
	L4	-0.8±0.15*	-0.75±0.17*	-0.4±0.13*	-0.4±0.13*	±	-0.6±0.15*	-0.8±0.12*	-0.73±0.11*
	L5	-0.74±0.12*	-1.4±0.24*	-0.42±0.15*	-0.21±0.2*	-0.38±0.36*	-0.47±0.14*	-0.51±0.14*	-0.18±0.18*
	L6	-0.9±0.2*	-0.41±0.24*	4.1±0.36*	-0.53±0.05*	-1.05±0.2*	-0.31±0.36*	-0.73±0.2*	-0.44±0.17*
	L7				1.65±0.51*				
	Adult	-0.1±0.1*	-0.84±0.32*	-0.62±0.2*	-0.22±0.22*	-0.77±0.18*	-0.45±0.26*	-0.56±0.2*	-0.82±0.08*±
Lti3	L1	-0.63±0.1*	0.43±0.54*	-0.68±0.9*	-0.68±0.9*	-0.31±0.42*	-0.59±0.31*	-0.58±0.19*	-0.41±0.22*
	L2	-0.99±0.2*	-0.03±0.4*	-0.55±0.3*	-0.55±0.3*	-0.19±0.17*	-0.85±0.3*	-0.2±0.2*	-0.4±0.3*
	L3	-0.64±0.15*	-0.61±0.19*	-0.84±0.4*	-0.84±0.4*	0.05±0.3*	-0.56±0.24*	-0.73±0.18*	-0.73±0.09
	L4	-0.75±0.18*	-0.2±0.19*	-0.16±0.16*	-0.16±0.16*	-0.22±0.27*	-0.6±0.2*	-0.26±0.34*	0.73±0.11*
	L5	-0.57±0.17*	-0.46±0.35*	0.08±0.9*	0.29±0.08*	-0.82±0.27*	-0.06±0.19*	-0.7±0.2*	-0.8±0.33*
	L6	-0.44±0.24*	-0.4±0.17*	-0.53±0.24*	-0.8±0.05*	-0.87±0.28*	-0.46±0.33*	-0.64±0.21*	-0.45±0.27*
	L7				0.9±0.2*				
	Adult	-0.06±0.08*	-0.13±0.21*	-0.54±0.24*	-0.2±0.34*	-0.86±0.09*	-0.47±0.41*	-0.11±0.19*	-0.82±0.08*

Notes: Each value of the table represents: mean and standard error. Asterisk (*), means that from the t-student test k is significantly different from 1. Legend: La: length of antenna. Labd: length of abdomen. Lai: length of the hind wing. Lcc: length of cephalic capsule. Lcu: width of cephalic capsule. Lcu1, Lcu2 and Lcu3: length of femur 1, length of femur 2, length of femur 3. Lel: length of elytra. Lpr: length of pronotum. Lti1, Lti2, Lti3: length of tibia 1, length of tibia 2, length of tibia 3. Lth: length of thorax. M: male. F: female. F6: female with six nymphal instars. F7: female with seven nymphal instars. L= nymphal instars. A= adult instars.

($k=1.66\pm 0.53$ for tibia 1 and $k=1.65\pm 0.51$ for tibia 2, $p<0,05$) (Table 5). Only 2 cases of isoallometries were noticed, mainly in the length of femur 3 in first nymphal instars females of *E. p. ibandana* ($k=0.99\pm 0.7$, $p>0,05$) (Table 5).

Discussion

This study showed that the growth model of the grasshopper species follows many types namely the power law, the Gompertz, the quadratic, the linear, the logarithm and the exponential models. This heterogeneous characteristic of the growth model is a common phenomenon in insects (Hirst and Forster 2013). However, for some authors, insect growth models are universal (West *et al.* 2001). These controversies could be due to the extreme sensibility of the larval growth ratio of insects to their living environment (temperature and humidity) and to the food quality accessible to them (Ayres and MacLean 1987; Esperk and Tammaru 2004). Thus, each larval instar might respond differently to these factors. These ideas might also explain the fact that the growth ratio of each organ between consecutive instars is not constant in each species and sex.

The power law model presented in this study as the best-fitted model for the studied grasshoppers would be a first observation in Acridomorpha group and might be linked to their feeding behaviour and developmental duration type. These grasshoppers share the same habitats and all are herbivorous and their developmental times are generally high in the first and last instars and low between those two instars. In general, being able to understand which model(s) best describe the growth curves, both empirically and ultimately mechanistically, is a real challenge (Hirst and Forster 2013). Only linear and exponential growth models were noted in Orthopteran populations (Von Bertalanffy 1951). At *Lucilia sericata* (Diptera: Calliphoridae) in South Korea, the logarithmic model was identified as the best fitting regression model (Shin *et al.* 2021). Invertebrate growth has often been modelled using exponential (Hirst and Bunker 2003; Nylin 1992; Hawkins 1986) or power mathematical form as the von Bertalanffy growth equation (von Bertalanffy 1957) and 73% growth of 58 marine invertebrates was best modelled by an exponential function (Hirst and Forster 2013). Additionally, sigmoidal functions, such as the von Bertalanffy growth equation, have been commonly applied to many vertebrates (Von Bertalanffy 1960; Kimura 1980; Zullinger 1984).

The limited amount of detailed information about the growth curves of insect larvae may partly result from technical difficulties (Tammaru and Esperk 2007). In particular, the existence of distinct larval instars makes larval growth curves complex, and any measurements of the instantaneous growth ratio should therefore explicitly consider the exact phase of the moulting cycle (Esperk and Tammaru

2004; Tammaru and Esperk 2007). Another drawback is the extreme sensitivity of larval growth ratios to environmental conditions, temperature and food quality, in particular (Tammaru and Esperk 2007).

The significant variation of growth ratios (growth coefficients) noticed in this study in each organ, between consecutive nymphal instars of each grasshopper species, reflect the positive covariance noticed in most Orthoptera by Whitman (2008) between most cuticular structures and the size of the individual of each species. Similar results were reported in earlier studies on various insects (Klingenberg and Zimmermann 1992). These results show that Dyar's rule does not apply in a strict sense to the growth of the material considered here. According to this law, each cuticular sclerite size increases in linear dimensions by a constant ratio (Dyar's coefficient or growth coefficient) at each moult. These different ratios noticed at the organ level of each species would reflect different growth strategies, and any change in the normal value of the growth coefficient for a species during a particular moult would represent an evolutionary adaptation of the growth pattern to environmental perturbations (Sehnal 1985, Whitman 2008).

The present study also showed that the allometric coefficient varied according to organs, sex, instars and species. The same conclusion has been obtained by Shingleton *et al.* (2007) with *Drosophila melanogaster* in USA and could be explained by the fact that this coefficient depends on environmental and physical conditions associated with increasing size (Whitman 2008).

The fact that hypoallometry model was the most adapted to describe the growth pattern of grasshoppers in this study suggests that in general, each organ of these grasshoppers experiences less relative growth compared to the overall growth of the organism. This result is close to that obtained in the Chinese dobsonfly (between male body size and ectoproct length (a male grasping structure)), in humans (between the skull and the human body) and in honey bees (abdomen, thorax, head, antenna, proboscis, pollen basket, maxillae, hind limb, fore wing and hind wing from the rainforest, Guinea savannah and the derived savannah zones of Nigeria) which exhibit negative allometric growth pattern (Cao *et al.* 2019; Bamidele 2021). This model may be dominant among grasshoppers and could be explained by the fact that the two parameters (size of organ and that of the whole body) are responding independently to what the growing insect is eating (Shingleton *et al.* 2007). Those grasshoppers are polyphytophagous and their organs respond differently to the variation of the characteristics factors of their habitat compared to the whole organism.

The few cases (7 cases) of hyperallometry noticed in the Width of cephalic capsule in the adult instar of *E. p.*

ibandana M; the Pronotum's length in *E. p. ibandana* F7 and the seventh nymphal instars; the lengths of tibia 1 and 2 in *E. p. ibandana* F6 and *E. p. ibandana* F7, in the sixth nymphal instars and the seventh nymphal instars, suggest the relatively faster growth of these organs compared to the overall growth of these grasshoppers. Similar results have been obtained by Cao et al. (2019) between male body size and mandible length of Chinese dobsonfly.

Only 2 cases of isoallometries growth were noticed, mainly in the length of femur 3 in first nymphal instars females of *E. p. ibandana*. Such isoallometric growth suggest that relative growth of a part of an organism is identical to the overall growth of that organism and would result in a dramatic mass-specific decrease in strength, because the maximum-force output of muscles scales proportionally to muscle cross-sectional area; but muscle cross-sectional area (and thus force output) increases proportionally less than mass, as size increases (Whitman 2008). However, in insects, there are usually isometric relationships between body size and other body parts (Cao et al. 2019).

In general, the nature of the nutrients circulating in the hemolymph affects the growth of insects. Morphology is correlated with the metabolic rate (Hirst et al. 2014, Glazier et al. 2015) and the type of allometric growth might depend on insect-specific assimilation rate. The specific assimilation varies during ontogenesis in many animals, resulting in greater absorptive capacity achieved by temporarily increasing organ size (Piersma et al. 1997, Hume et al. 2002). Increasing specific assimilation implies that organs do not develop isomorphically. Consistent specific assimilation can be expected if organs related to nutrient uptake grow isomorphically with body size (Maino and Kearney 2015). The use of these nutrients is also dependent on circulating hormone peaks (Joly 1968, David et al. 1999). For example, the water metabolism of locusts is dependent on diuretic (AVP-like Insect Diuretic hormone, peptidergic in nature) and antidiuretics (Neuroparsine, Chloride Transport Stimulating Hormone, antidiuretic hormone) hormones. Neurosparin is also involved in energy metabolism and phase polymorphism of Locusts (Proux, 1991).

The insulin / IGF (« Insulin-like Growth factor or Somatomedin ») signaling which play a key role in the longevity, diabetes, and the regulation of the size of cells, organs and the whole body could equally have a central role in the regulation of the organism shape (Nijhout et Callier 2015). In *D. melanogaster* nutrition poor in amino acids can be directly detected

be dividing cells, causing a reduction of the insulin signalling pathway in these cells and then a reduction of the growth ratio (Neufeld 2003, Oldham et al. 2000, Goberdhan and Wilson 2002). Then, the allometric variation could be due to the organ 'specific changes in response to the hormones (Shingleton et al. 2007).

Most authors including Shingleton et al. (2007) draw their conclusions on the type of allometry (Hyper-, Iso- and Hypoallometries) based only on a simple comparison of the coefficient k to 1 (without statistical test). The application of a statistical test as in this study to make such comparison facilitated the interpretation of this coefficient and help to draw robust conclusions. However, the ease of calculation and interpretation of the allometric coefficient made the model generally useful, but there is some question about the capacity of a power function of the type $y = ax^b$ to describe relative growth (Bervian et al. 2006).

Conclusion

The growth models of *A. acutipennis*, *E. p. ibandana*, *P. vignaudii* and *T. ferruginea* are not unique. Under laboratory conditions, those grasshoppers have been represented by six types of growth models: power, gompertz, quadratic, linear, logarithm and exponential. The power law model was the best-fitted one in all the grasshopper species. The growth ratios of elytra length and hind wing length were high and varied between 2 to 4 except for *E. p. ibandana* F7 where the multiplication factors from nymphal instars 6 to 7 were close to 1; meaning that the wings' size lowly changed between the two stages. For each organ, a disproportionate variation in the allometric coefficient from one instar to another was observed. These findings will go a long way to improve the knowledge of the characteristics of acridians inhabiting forest agroecosystems.

Ontogeny is affected by both genetic and environmental factors that operate through complex molecular and physiological mechanisms. Anyway, we are still a long way from understanding the mechanisms that underlie nymphal development scaling patterns of insects.

Acknowledgements

We thank Pr Ajeagah Gideon of the University of Yaoundé 1 (Cameroon), who revised this manuscript as a native English speaker. We also thank all the co-authors of Kekeunou et al. (2020), Djomang Nkwala et al. (2019), Kekeunou et al. (2018), Kekeunou et al. (2015) and members of the Plant Protection Service, Cameroon (2PC) who are not co-authors in this paper, for data collection in laboratory.

References

- Abrams PA, Leimar O, Nylin S, Wiklund C. The effect of flexible growth rates on optimal sizes and development times in seasonal environment. *Am. Nat.* 1996; 147: 381-395.
- Arnhold E. Easy Nonlinear Model, version 5.0: Fit and plot some nonlinear models. Repository CRAN; 2017. <https://cran.r-project.org>
- Ayres MP, MacLean SF. Molt as a component of insect development, *Gallerucella sagittariae* (Chrysomelidae) and *Epirrita autumnata* (Geometridae). *Oikos* 1987; 48: 273-279.
- Bamidele AJ, Idowu BA, Ademolu OK, Osipitan AA. Allometric studies on *Apis mellifera adansonii* workers from three ecological zones of Nigeria. *Bulletin of Insectology* 2021; 74 (1): 19-26.
- Berger D, Walters R, Gotthard K. What keeps insects small? Size dependent predation on two species of butterfly larvae. *Evol. Ecol.* 2006; 20: 575-589.
- Bervian G, Fontoura NF, Haimovici M. Statistical model of variable allometric growth: otolith growth in *Micropogonias furnieri* (Actinopterygii, Sciaenidae). *Journal of Fish Biology* 2006; 68: 196-208.
- Blossman-Myer B, Burggren WW. Metabolic Allometry during Development and Metamorphosis of the Silkworm *Bombyx mori*: Analyses, Patterns, and Mechanisms. *Phys. Bioc. Zool.* 2010; 83 (2): 215-231.
- Canard A, Poinso D. Quelques méthodes statistiques typiques de l'étude des populations et des peuplements par la méthode des quadrats. Fiche technique. Université de Renne 1 (France) 2004 ; 34 p.
- Cao C, Yu P, Hayashi F. Allometry and morphological trait relationship in the sexually dimorphic Chinese dobsonfly, *Acanthacorydalis asiatica* (Wood-Mason, 1884) (Megaloptera, Corydalidae). *ZooKeys* 2019; 854: 119-129.
- David L, Stern, Douglas J, Emlen. The developmental basis for allometry in insects. *Development* 1999; 126: 1091-1101.
- Day T, Taylor PD. Von Bertalanffy's growth equation should not be used to model age and size at maturity. *Am. Nat.* 1997; 149: 381-393.
- De Grégorio R. *Zonocerus variegatus* (Orthoptera, Pyrgomorphidae) : caractéristiques morphologiques et biométriques des larves des populations des saisons sèche et humide. *Bul. Soc. Hist. Nat. Toul.* 1987 ; 123: 29-44.
- Default B. Biométrie des types de Caelifères de France (Orthoptera). 1. Définition des paramètres mesurés. 2. Mesurations chez les Tridactylidae, Tetrigidae, Pyrgomorphidae, Pamphagidae et Acrididae Calliptaminae. *Mat. Orth. Ent.* 2012 ; 17 :21-56.
- Djomnang Nkwala AL, Simo Talla F, Oumarou Ngoute C, Kekeunou S, Wandji AC, Mbadjoun Nzike M, Simeu Noutchom A, Mpoame Mbida. Morphology, development, and reproduction of *Eyprepocnemis plorans ibandana* Giglio-Tos, 1907 (Orthoptera, Acrididae) in south Cameroon rainforests. *J. Orth. Res.* 2019; 28 (2): 145-154.
- Enquist BJ, West GB, Charnov EL, Brown JH. Allometric scaling of production and life-history variation in vascular plants. *Nature* 1999; 401: 907-911.
- Esperk T, Tammaru T. Does the 'investment principle' model explain molting strategies in Lepidopteran larvae? *Phys. Ento.* 2004; 29: 56-66.
- Fenchel T. Intrinsic rate of natural increase: the relationship with body size. *Oecologia* 1974; 14: 317-326.
- Glazier DS, Hirst AG, Atkinson D. Shape shifting predicts ontogenetic changes in metabolic scaling in diverse aquatic invertebrates. *Proc. R. Soc. B.* 2015; 282: 1-9. doi:10.1098/rspb.2014.2302-34.
- Goberdhan DC, Wilson C. Insulin receptor-mediated organ overgrowth in *Drosophila* is not restricted by body size. *Dev. Genes Evol.* 2002; 212 (4): 196-202.
- Gockowski J, Tonye J, Baker D, Legg C, Weise S, Tchienkoua M, Ndoumbe M, Tiki Manga T, Fouaguégué. Characterization and diagnosis of farming systems in the forest margins benchmark of southern Cameroun. International Institute of Tropical Agriculture 2004; Social Sciences Working Paper Series 1.
- Gotthard K. Growth strategies and optimal body size in temperate Parargini butterflies. *Int. Comp. Biol.* 2004; 44: 471-479.
- Hawkins CP. Variation in individual growth rates and population densities of ephemeropterid mayflies. *Ecology* 1986; 67: 1384-1395.
- Hirst A, Forster J. When growth models are not universal: evidence from marine invertebrates. *Proc. Royal. Soc. B.* 2013; 280: 15-46.
- Hirst AG, Bunker AJ. Growth of marine planktonic copepods: global rates and patterns in relation to chlorophyll a, temperature, and body weight. *Limnol Oceanogr* 2003; 48: 1988-2010.
- Hirst AG, Glazier DS, Atkinson D. Body shape shifting during growth permits tests that distinguish between competing geometric theories of metabolic scaling. *Ecol. Lett.* 2014; 17: 1274-1281. doi:10.1111/ele.12334
- Honek A. Intraspecific variation in body size and fecundity in insects: a general relationship. *Oikos* 1993; 66: 483-492.
- Hume D, Beiglböck C, Ruf T, Frey-Roos F, Bruns U, Arnold W. Seasonal changes in morphology and function of the gastrointestinal tract of free-living alpine marmots (*Marmota marmota*). *J. Comp. Physiol. B.* 2002; 172 : 197-207. doi:10.1007/s00360-001-0240-1
- Joly P. *Endocrinologie des Insectes*. Masson & Cie 1968 ; 344p.
- Kekeunou S, Mbadjoun-Nziké M, Wandji AC, Baleba BS, Djomnang-nkwala AL, Simeu-Noutchom A, Oumarou-Ngoute C, Um-Nyobe PCA, Guiadem-Simo L, Ntonga Akono P. Morphology, development and reproduction of *Atractomorpha acutipennis* (Guérin-Méneville, 1844) (Orthoptera: Pyrgomorphidae). *Trop. Zool.* 2020 ; 33 (3): 97-112. doi:10.4081/tz.2020.81
- Kekeunou S, Mbeng D, Oumarou Ngoute C, Wandji AC. Morphology, development and reproduction of *Pyrgomorpha vignaudii* (Orthoptera: Pyrgomorphidae). *Entomol. Res.* 2015; 45: 58-70.
- Kekeunou S, Wandji AC, Oumarou Ngoute C. Morphology, post-embryonic development and reproduction of *Taphronota ferruginea* (Fabricius, 1781) (Orthoptera: Pyrgomorphidae). *Trop. Zool.* 2018; 31 (2): 68-84.
- Kimura DK. Likelihood methods for the von Bertalanffy growth curve. *Fish. Bul.* 1980; 77: 765-776.
- Kjørboe T, Hirst AG. Optimal development time in pelagic copepods. *Mar. Ecol. Prog. Ser.* 2008; 367: 15-22.
- Klingenberg C P, Zimmermann M. Dyar's rule and multivariate allometric growth in nine species of waterstriders (Heteroptera: Gerridae). *J. Zool. Lond.* 1992; 227: 453-464.
- La Barbera M. Analyzing body size as a factor in ecology and evolution. *Ann. Rev. Ecol. Syst.* 1989; 20: 97-117.
- Lee L, Atkinson D, Hirst AG, Cornell SJ. A new framework for growth curve fitting based on the von Bertalanffy Growth Function. *Sci. Rep.* 2020; 10 (7953): 34p. <https://doi.org/10.1038/s41598-020-64839-y>
- Maino JL, Kearney MR. Testing mechanistic models of growth in insects. *Proc. R. Soc. B.* 2015; 282: 1973. <http://dx.doi.org/10.1098/rspb.2015.1973>
- Neufeld TP. Body building: regulation of shape and size by PI3K/TOR signaling during development. *Mech. Dev.* 2003; 120: 1283-1296.
- Nijhout HF, Callier V. Developmental Mechanisms of Body Size and Wing-Body Scaling in Insects. *Ann. Rev. Entomol.* 2015; 60: 141-56.
- Nylin S. Seasonal plasticity in life history traits: growth and development in *Polygonia calbum* (Lepidoptera: Nymphalidae). *Biol. J. Linn. Soc.* 1992; 47: 301-323.
- Oldham S, Montagne J, Radimerski T, Thomas G, Hafen E. Genetic and biochemical characterization of dTOR, the *Drosophila* homolog of the target of rapamycin. *Genes Dev* 2000; 14: 2689-2694.
- Piersma T, Lindström A. Rapid reversible changes in organ size as a component of adaptive behaviour. *Trends Ecol. Evol.* 1997; 12: 134-138. doi:10.1016/S0169-5347(97)01003-3
- Proux J. La régulation hormonale du métabolisme hydrique chez les criquets grégaires. In: Abdourrahmane Essaid (Ed): *La lutte antiacridienne*. Aupelf-UREF, John Libbey Eurotext 1991 ; pp

129-134.

Ricklefs RE. Is rate of ontogenetic growth constrained by resource supply or tissue growth potential? A comment on West *et al*'s model. *Funct. Ecol.* 2003; 17: 384-393.

Sehnal F. Growth and life cycles, pp. 2-86. In: Kerkut G.A., Gilbert L.I. (Eds). *Comprehensive Insect Physiology, Biochemistry and Pharmacology* 1985; Vol 2. Pergamon Press, Oxford.

Shin SE, Park JH, Jeong SJ, Park SH. The Growth Model of Forensically Important *Lucilia sericata* (Meigen) (Diptera: Calliphoridae) in South Korea. *Insects* 2021; 120423. <https://doi.org/10.3390>

Shingleton AW, Frankino WA, Flatt T, Nijhout HF, Emlen DJ. Size and shape: the developmental regulation of static allometry in insects. *BioEssays* 2007; 29: 536-548.

Stoica P Selen Y. Model-order selection: a review of information criterion rules. *IEEE Signal Processing Magazine* (July), 2004: 36-47.

Stoner MP. Fitting the Exponential Function and the Gompertz Function by the Method of Least Squares. *Journal of the American Statistical Association* 1941; 36 (216): 515-518.

Suchel JB. Les climats du Cameroun. Les climats du Cameroun. Thèse de doctorat d'état. Université de Bordeaux III, 2 tomes, 1987. 186 p.

Tammaru T, Esperk T. Growth allometry of immature insects: larvae do not grow exponentially. *Funct. Ecol.* 2007; 21: 1099-1105.

Tammaru T, Esperk T, Castellanos I. No evidence for costs of being large in females of *Orgyia* spp. (Lepidoptera, Lymantriidae): larger is always better. *Oecologia* 2002; 133: 430-438.

Von Bertalanffy L. Metabolic types and growth types. *Am. Nat.* 1951; 85 (821): 111-117.

Von Bertalanffy L. Quantitative laws in metabolism and growth. *Quart. Rev. Biol.* 1957; 32: 217-231.

Von Bertalanffy L. Principles and theory of growth. In *Fundamental aspects of normal and malignant growth* (ed. WW Nowinski) New York, NY: Elsevier, 1960: 137-259.

West GB, Brown JH, Enquist BJ. A general model for ontogenetic growth. *Nature* 2001; 413: 628-631.

Withman DW. The significance of body size in the Orthoptera: a review. *J. Orth. Res.* 2008; 17(2): 117-134.

Yang Y, Joern A. Gut size changes in relation to variable food quality and body size in grasshoppers. *Functional Ecology* 1994; 8: 36-45.

Zullinger EM. Fitting sigmoidal equations to mammalian growth curves. *J. Mammal.* 1984; 65: 607-636.

Mini Curriculum and Author's Contribution

1. Sévilor Kekeunou, PhD; Professor. Contribution: designed the study, analysed the data, interpreted and discussed the results, and contributed to the text. ORCID: 0000-0002-9058-100X

2. Alfiery Laurel Djomnang-Nkwala, PhD. Contribution: designed the study, analysed the data, interpreted and discussed the results, and contributed to the text. ORCID: 0000-0001-7887-9535

3. Alain Christel Wandji, PhD. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0002-4394-9502

4. Richard Akwandjoh Seino, PhD; Associate Professor. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0003-4379-9871

5. Marcelle Mbadjoun Nzike, PhD student. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0001-5327-1638

6. Charly Oumarou Ngoute, PhD. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0002-1257-3573

7. Alain Simeu Noutchom, PhD. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0002-5386-9450

8. Franklin Simo Talla, PhD student. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0002-2607-9648

9. Donatus Mbeng, PhD. Contribution: interpreted and discussed the results, and contributed to the text. ORCID: 0000-0002-8806-9816

10. Mpoame Mbida, PhD; Professor. Contribution: interpreted and discussed the results, and contributed to the text.

Received: March 20, 2023

Accepted: May 5, 2023

Corresponding author

Sévilor Kekeunou

E-mail: skekeunou@gmail.com